

Supporting Information

Precise Synthesis of Bottlebrush Block Copolymers from ω -End-Norbornyl Polystyrene and Poly(4-*tert*-butoxystyrene) via Living Anionic Polymerization and Ring-Opening Metathesis Polymerization

Yong-Guen Yu,^{†,‡} Chang-Geun Chae,^{†,‡} Myung-Jin Kim,^{†,‡} Ho-Bin Seo,^{†,‡} Robert H. Grubbs,^{‡,§} and Jae-Suk Lee^{*,†,‡}

[†]School of Materials Science and Engineering and [‡]Grubbs Center for Polymers and Catalysis, Gwangju Institute of Science and Technology (GIST), 123 Cheomdangwagi-ro, Buk-gu, Gwangju 61005, Republic of Korea.

[§]Division of Chemistry and Chemical Engineering, California Institute of Technology (Caltech), Pasadena, California 91125, United States.

E-mail: jslee@gist.ac.kr; Tel: (+82)-62-715-2306.

Contents

1. ¹H and ¹³C NMR spectra of chemical compounds
2. Physical data of macromonomer
3. Additional SEM images and photographs of films
3. Additional kinetic profile on ROMP of NPSt in THF at room temperature ([NPSt]₀ = 0.05 M)

1. ^1H and ^{13}C NMR spectra of chemical compounds




Figure S1. ^1H NMR spectrum of 2,2,5,5-tetramethyl-1-(3-chloropropyl)-1-aza-2,5-disilacyclopentane (T1) in CDCl_3 (400 MHz).




Figure S2. ^{13}C NMR spectrum of 2,2,5,5-tetramethyl-1-(3-chloropropyl)-1-aza-2,5-disilacyclopentane (T1) in CDCl_3 (100 MHz).




Figure S3. ¹H NMR spectrum of *N*-(carboxylhexanoyl)-*cis*-norbornene-*exo*-2,3-dicarboxiimide in CDCl₃ (400 MHz).




Figure S4. ¹³C NMR spectrum of *N*-(carboxylhexanoyl)-*cis*-norbornene-*exo*-2,3-dicarboxiimide in CDCl₃ (100 MHz).




Figure S5. ^1H NMR spectrum of *N*-(perfluorophenyl hexanoate)-*cis*-norbornene-*exo*-2,3-dicarboxiimide (T2) in CDCl_3 (400 MHz).




Figure S6. ^{13}C NMR spectrum of *N*-(perfluorophenyl hexanoate)-*cis*-norbornene-*exo*-2,3-dicarboxiimide (T2) in CDCl_3 (100 MHz).




Figure S7. ^9F NMR spectrum of *N*-(perfluorophenyl hexanoate)-*cis*-norbornene-*exo*-2,3-dicarboxiimide (T2) in CDCl_3 (376 MHz).




Figure S8. ^1H NMR spectrum of PSt-NH_2 in CDCl_3 (400 MHz).




Figure S9. ¹H NMR spectrum of PtBOS-NH₂ in CDCl₃ (400 MHz).




Figure S10. ¹H NMR spectrum of NPSt in CDCl₃ (400 MHz).




Figure S11. ^1H NMR spectrum of NPtBOS in CDCl_3 (400 MHz).




Figure S12. ^1H NMR spectrum of P(NB-g-St)₆₃₀ in CDCl_3 (400 MHz).




Figure S13. ^1H NMR spectrum of $\text{P}(\text{NB-g-tBOS})_{154}$ in CDCl_3 (400 MHz).




Figure S14. ^1H NMR spectrum of $\text{P}(\text{NB-g-St})_{151}\text{-b-P}(\text{NB-g-tBOS})_{140}$ in CDCl_3 (400 MHz).

2. Physical data of macromonomers

Table S1. Characteristics of ω -norbornyl macromonomers

entry	MM	$M_{n,\text{NMR}}$ (kDa)	$M_{n,\text{obsd}}^a$ (kDa)	\bar{D}^a	DP_n
1	NPSt	2.76	2.69	1.14	24
2	NPSt	3.23	3.37	1.07	28
3	NPtBOS	2.79	2.43	1.06	14
4	NPtBOS	3.13	3.01	1.06	16

^a $M_{n,\text{obsd}}$ and \bar{D} were obtained by SEC calibration using polystyrene standard in THF containing 2% trimethylamine as the eluents at 40 °C.




Figure S15. GPC curves of the macromonomers of (a) NPSt and (b) NPtBOS

3. Additional SEM images and photographs of films of Bottlebrush block copolymers




Figure S15. Cross-sectional SEM image of P(NB-g-St)-*b*-P(NB-g-*t*BOS) with $M_w = 148$ kDa and $DP_w = 52$ (Table 2, entry 1).




Figure S16. Cross-sectional SEM images and photographs of photonic films of P(NB-g-St)-*b*-P(NB-g-*t*BOS) with (a,b) $M_w = 296$ kDa (Table 2, entry 2), (c,d) $M_w = 830$ kDa (Table 2, entry 3), and (e,f) $M_w = 1331$ kDa (Table 2, entry 4).

**4. Additional kinetic profile on ROMP of NPSt in THF at room temperature
([NPSt]₀ = 0.05 M)**




Figure S17. SEC traces of P(NB-g-St) aliquots with [NPSt]₀/[G3]₀ = 650 withdrawn from mixture during ROMP at time intervals (Table1, entry 3).